

2018 RESEARCH REPORT**Principle Investigator:****Carmela M. Buono, PhD Student**

Dr. Kirsten Prior, Faculty Advisor

Department of Biological Sciences
 Binghamton University, State University of New York
 PO Box 6000, Vestal, New York 13902
 Phone (cell): (732) 425-2666
 Email: cbuono1@binghamton.edu

Project Title: Historical land-use impact of ant-mediated seed dispersal in northeastern deciduous forests

Location: *Cornell Botanical Gardens (Bald Hill, Carter Creek, Fall Creek Valley North and South, and Slaterville 600)*

I. Study Purpose and Objectives

As humans increasingly develop and change the surface of the earth, large impacts are being made to natural communities such as habitat loss and a reduction in biodiversity. One of the largest anthropogenic contributors to the disruption of natural ecosystems is *land-use change* [1]. Historical changes in land-use can leave legacy effects such as, habitat fragmentation, changes in soil chemistry, and altered species composition [2–4]. With the additional threat of anthropogenic climate change, it is becoming increasingly important to understand how multiple anthropogenic stressors are impacting our ecosystems so that we may better conserve and restore them. In order to implement effective management and restoration practices, we first must understand if ecological systems can recover from the impacts of land-use change. By understanding if and in what contexts systems can recover from land-use change, then we can more appropriately apply management and restoration practices. This project focuses on one form of anthropogenic impacts, *land-use change history*, and how they are impacting ecosystem biodiversity and function. This project addresses the research questions below and sections II.-III. will detail preliminary data from summer 2018 as well as future and continued work.

Human alteration of landscapes is the major cause of biodiversity loss and of the disruption of ecosystem functions [5]. The majority of forests in northeastern North America have at one time been cleared for agricultural or timber use but recently these fragments of previously cleared land have been abandoned and allowed to passively regenerate [6]. As a result, contemporary forests are composed largely of “secondary” or previously cleared forest, with less than 1% of “primary” forests remaining [6–9]. Given that much of our natural lands have already been converted for food production and living space, with continuing pressure [5], a key question for ecologists tasked with conserving and restoring natural systems is whether these systems can recover from large-scale land-use change disturbances. To this end, we can look to natural areas that have passively recovered after previous land-use and measure ecological responses to better understand how we can successfully restore ecosystems.

In northeastern forests, entire forest compositions were altered when forests were cleared. In secondary forests, it has been observed the tree and shrub layers have successfully recovered passively [6], while understory plant communities, which represent the greatest plant diversity in

forests, seem to be greatly reduced in richness and abundance [6,10]. Particularly absent in these understory communities are myrmecochorous plants [11], which are plants specialized for seed dispersal by ants (i.e., their seeds have a lipid-rich appendage, “elaiosome”) [8]. Myrmecochores are particularly rich in eastern deciduous forests, making up 35-40% of understory species, and including many well-known species, such as *Asarum canadense*, *Sanguinaria canadensis*, *Uvularia perfoliata*, *Anemone acutiloba*, *Trillium* sp., *Viola* sp., and *Dicentra* spp. [12–14]. Ants are attracted to and use the elaiosome to carry the seeds to their nests where they remove the elaiosome and feed it to their larvae, after which they “plant” the intact seed outside of their nests. This is a mutualistic interaction, in which ants receive a food reward while myrmecochores benefit by having their seeds dispersed away from conspecifics and protected from seed predators [15,16]. In eastern forests, ants in the genus *Aphaenogaster* are responsible for the majority of seed dispersal and are often referred to as “keystone dispersers” [17]. Seed dispersal by ants is an essential ecosystem function for understory plants in deciduous forests, yet few studies have examined how land-use change affects this interaction, specifically across different regions in the northeast [17–19]. In order to conserve and work to restore the understory forest plant community in secondary forests, we need to better understand how seed dispersal is impacted, and if and how we may need to also restore or enhance this key function in secondary forests [17]. **In my research, I am investigating if ant-mediated seed dispersal is intact in secondary forests by conducting a large-scale natural experiment, comparing myrmecochores, ants, ant habitat, and dispersal rates in primary and secondary forests in across the northeast.**

Main Question:

Does land-use history alter ecological function in secondary forests? I will examine how land-use history affects an important ecosystem function, ant-mediated seed dispersal in eastern deciduous forests. In particular, I will examine if land-use affects the presence and strength of seed dispersal function by conducting observational surveys of vegetative communities, ant mutualist presence and abundance, and presence of seed dispersal function in paired primary and secondary forests in the northeast.

A1. PATTERNS IN MYRMECOCHOUS PLANT COMMUNITIES AND ANT HABITAT *by surveying the abundance and diversity of forest understory plants, including myrmecochores and the type and abundance of potential nesting sites.*

A2. PATTERNS IN ANT-MEDIATED SEED DISPERSAL *by surveying the identity and abundance of seed-dispersing ants and measuring mutualist function (seed dispersal rates).*

To this end, I have identified 20 primary and secondary forest sites in three regions (A: New Jersey, eastern New York; B: northern PA; C: central NY). In 17 sites, I conducted vegetative surveys (**including Bald Hill, Carter Creek, Fall Creek Valley North and South, and Slaterville 600**), and I am still analyzing this data. In 11 sites, I conducted seed dispersal trials and pitfall traps (**including Bald Hill and Slaterville 600**), and I am finding that that seed dispersal is lower and more variable in secondary forests as well as the abundance of the

keystone seed-dispersing ants (ants in the genus *Aphaenogaster*). This summer I will finish the invertebrate surveys at the remaining 6 sites leftover from summer 2018. In addition, I will be adding the final 3 sites to reach a total of 20 forests across the northeast where I will complete the same vegetative and invertebrate surveys.

II. Methods

This past summer, I observed the interactions of ant mutualists on the native herbaceous community. I chose to work at **Bald Hill, Carter Creek, Fall Creek Valley North and South, and Slaterville 600** due to it containing secondary and old-growth forest habitat where the key mutualism for this project is found. To do this, I conducted i) vegetative surveys, ii) ant habitat surveys, iii) ant community surveys, and iv) seed dispersal rate measurements. These surveys consisted of three 50-meter semi-permanent belt transects that will be surveyed for another summer (Fig. 4). The transects are located in the center of the forest (50 meters or more away from an edge). Each transect includes 20 5*5m survey plots, occurring on either side of the transect (5 as “plant” plots, and 5 as “invertebrate” plots, 15 of each plot per site). The surveys are observational and did *not* include any damaging of specimens.

To mark the plots for semi-permanent use, 1-foot 1-inch-wide PVC pipe were installed at each end of the transects. These posts are blue in color and look like (Fig. 3). Each PVC post has contact and identification information laminated and attached to the post.

i). The vegetative surveys of the herbaceous layer compare diversity, cover, and composition of vegetative species in the 5x5 “plant” plots. I measured the presence and cover of understory plants (including myrmecochores) in 4, 1 m² quadrats within plant plots. In each plot, I also measured shrub and tree composition and cover, along with potential correlates that could influence understory plants, such as soil temperature, soil moisture, soil pH, patch size, and light availability, and ground cover. The vegetative surveys are observational and did not include any damaging of specimens. Exact measurements in the vegetative plots can be found in Table 1.

ii). In a subset of plots (at only **Bald Hill and Slaterville 600**), I sampled the ant community by placing pitfall traps (7 cm deep, 9 cm wide, and 9 oz) out for 24 hours. Two pitfall traps were in the “invertebrate” plots (total site n = 30) and placed flush with ground level. A combination of soap and water were placed into these cups and left out for 24 hours. A wire mesh filter was placed on top of each of these containers to eliminate the chances of small rodents or salamanders accidentally being caught. After 24 hours, the cups were collected, and the contents of the containers were emptied and later identified. These pitfall traps were left out for the 24 hours on a dry day and collected (cups and all other materials) and removed from the ground immediately (leaving no trace).

iv). Finally, to test the ability of the ants to disperse seeds, seed depots with 8 native plant seeds were placed out with a mesh covering (to exclude rodents). These depots were left out for a total of 24 hours on a dry day and the presence or absence of seeds were recorded the next day. On the final day, all materials were removed and from the ground (leaving no trace). Seeds were from native *Asarum canadense* and are not viable. From completed iv), I am already finding

interesting results - consistent seed removal in primary forests and low and variable removal in secondary forests (Fig. 2).

The primary goal of this summer was to gather baseline data of the biotic interactions that I will be measuring between primary and secondary forests. In addition, I used this opportunity to indicate the locations for my vegetative and ant surveys so that I can return to them for the next summer to collect additional data. This work is part of a PhD dissertation, in which I will pair this natural experiment with a manipulative experiment.

III. Preliminary Results

Out of 20 sites, I have completed 17 of (Obj.1) and 11 of (Obj.2 & Obj. 3). This upcoming field season, I plan to finish plant and ant surveys in the summer of 2019. I will perform linear or generalized linear models and AIC (Akaike's Information Criterion) to test if region, land-use history, and other measured variables affects plant, ant, and habitat response variables. From completed iv), I am already finding interesting results - consistent seed removal in primary forests and low and variable removal in secondary forests (Fig.2). **This work is part of a PhD dissertation**, in which I will pair this natural experiment with a manipulative experiment.

Table 1. Vegetative species identified (to lowest taxonomic group possible) during vegetative surveys at Cornell Botanic Gardens Property, summer 2018. We are currently identifying the herbaceous species.

Genus	Species	Genus	Species
Canopy Layer Bald Hill		Shrub Layer Bald Hill	
<i>Acer</i>	<i>rubrum</i>	<i>Acer</i>	<i>pensylvanicum</i>
<i>Betula</i>	<i>lenta</i>	<i>Hamamelis</i>	<i>virginiana</i>
<i>Carpinus</i>	<i>caroliniana</i>	<i>Kalmia</i>	<i>latifolia</i>
<i>Crataegus</i>	<i>pruinosa</i>	<i>Lindera</i>	<i>benzoin</i>
<i>Fagus</i>	<i>grandifolia</i>	<i>Lonicera sp.</i>	
<i>Fraxinus</i>	<i>americana</i>	<i>Vaccinium sp.</i>	
<i>Hamamelis</i>	<i>virginiana</i>	Shrub Layer Carter Creek	
<i>Nyssa</i>	<i>sylvatica</i>	<i>Acer</i>	<i>pensylvanicum</i>
<i>Ostrya</i>	<i>virginiana</i>	<i>Berberis</i>	<i>thunbergii</i>
<i>Quercus</i>	<i>alba</i>	<i>Rosa</i>	<i>multiflora</i>
<i>Quercus</i>	<i>motana</i>	Shrub Layer Fall Valley Creek	
<i>Quercus</i>	<i>rubra</i>	<i>Berberis</i>	<i>thunbergii</i>
<i>Tsuga</i>	<i>canadensis</i>	<i>Lindera</i>	<i>benzoin</i>
Canopy Layer Carter Creek		<i>Lonicera</i>	<i>morrowii</i>
<i>Acer</i>	<i>rubrum</i>	<i>Rubus</i>	<i>occidentalis</i>
<i>Acer</i>	<i>saccharum</i>	Shrub Layer Slaterville 600	
<i>Betula</i>	<i>lenta</i>	<i>Acer</i>	<i>pensylvanicum</i>
<i>Carpinus</i>	<i>caroliniana</i>		
<i>Fagus</i>	<i>grandifolia</i>		
<i>Fraxinus</i>	<i>americana</i>		
<i>Ostrya</i>	<i>virginiana</i>		
<i>Tilia</i>	<i>americana</i>		
<i>Tsuga</i>	<i>canadensis</i>		
Canopy Layer Fall Creek Valley			
<i>Fagus</i>	<i>grandifolia</i>		
<i>Acer</i>	<i>saccharum</i>		
<i>Fraxinus</i>	<i>americana</i>		
<i>Ostrya</i>	<i>virginiana</i>		
<i>Prunus</i>	<i>serotina</i>		
<i>Tsuga</i>	<i>canadensis</i>		
<i>Tilia</i>	<i>americana</i>		
<i>Acer</i>	<i>rubrum</i>		
<i>Carya</i>	<i>glabra</i>		
Canopy Layer Slaterville 600			
<i>Acer</i>	<i>saccharum</i>		
<i>Carpinus</i>	<i>caroliniana</i>		
<i>Fagus</i>	<i>grandifolia</i>		
<i>Fraxinus</i>	<i>americana</i>		
<i>Ostrya</i>	<i>virginiana</i>		
<i>Tilia</i>	<i>americana</i>		


Table 2. Insect Orders identified from the pitfall traps implemented for the ant abundance surveys at Cornell Botanic Gardens Property, Summer 2018. We are currently identifying the ants to species.

Code	Order
Slaterville 600	
ANA	<i>ANNELIDA</i>
ARA	<i>ARANEAE</i>
CHI	<i>CHILOPODA</i>
COLE	<i>COLEOPTERA</i>
COLL	<i>COLLEMBOLA</i>
DIPT	<i>DIPTERA</i>
ISOP	<i>ISOPODA</i>
LEP	<i>LEPIDOPTERA</i>
OPI	<i>OPILIONES</i>
Bald Hill	
ACA	<i>ACARI</i>
ANA	<i>ANNELIDA</i>
ARA	<i>ARANEAE</i>
CHI	<i>CHILOPODA</i>
COLE	<i>COLEOPTERA</i>
COLL	<i>COLLEMBOLA</i>
DIPL	<i>DIPLOPODA</i>
DIPT	<i>DIPTERA</i>
GAST	<i>STYLOMMATOPHORA</i>
HYM	<i>HYMENOPTERA</i>
ISOP	<i>ISOPODA</i>
LEP	<i>LEPIDOPTERA</i>
OPI	<i>OPILIONES</i>


Table 3. GPS locations of transect plots at Cornell Botanic Gardens Property, Summer 2018

Marker Code	Lat	Long	Marker Code	Lat	Long
Slaterville 600			Bald Hill		
CN1PAL	N 42.422233	W 76.326033	CN1SAL	N 42.355283	W 76.382133
CN1PAR	N 42.421917	W 76.325517	CN1SAR	N 42.354883	W 76.381683
CN1PBL	N 42.421817	W 76.327200	CN1SBL	N 42.354700	W 76.381383
CN1PBR	N 42.421350	W 76.326683	CN1SBR	N 42.354283	W 76.380983
CN1PCL	N 42.421033	W 76.328033	CN1SCL	N 42.353850	W 76.381050
CN1PCR	N 42.420700	W 76.327583	CN1SCR	N 42.353400	W 76.381000
Carter Creek					
CN2PAL	N 42.333683	W 76.664433	CN2SAL	N 42.337883	W 76.662567
CN2PAR	N 42.333733	W 76.665100	CN2SAR	N 42.337950	W 76.661967
CN2PBL	N 42.334550	W 76.665367	CN2SBL	N 42.338450	W 76.662367
CN2PBR	N 42.334483	W 76.665950	CN2SBR	N 42.338167	W 76.662883
CN2PCL	N 42.335000	W 76.665567	CN2SCL	N 42.337050	W 76.661583
CN2PCR	N 42.334933	W 76.666167	CN2SCR	N 42.337000	W 76.662183
CN3PAL	N 42.455083	W 76.451233	CN3SAL	N 42.458333	W 76.449417
CN3PAR	N 42.455033	W 76.450700	CN3SAR	N 42.457933	W 76.449083
CN3PBL	N 42.455361	W 76.450528	CN3SBL	N 42.458783	W 76.449100
CN3PBR	N 42.454867	W 76.450367	CN3SBR	N 42.458417	W 76.448667
CN3PCL	N 42.455100	W 76.450367	CN3SCL	N 42.458600	W 76.449200
CN3PCR	N 42.455467	W 76.450067	CN3SCR	N 42.458222	W 76.448778

A.



B.






Figure 1. Map of survey regions on Cornell Botanic Gardens Property, Summer 2018 A. Bald Hill, B. Carter Creek, C. Fall Creek Valley, and D. Slaterville 600.




Figure 2. Boxplots of the portion of seeds removed from 5 primary and 6 secondary forest sites in the northeast. ($p= 0.0393$)



Figure 3. An image of the .5-foot-tall and 1-inch wide PVC posts that will be used to identify the ends of the study transects. In addition, these PVC posts are blue and have contact and identification information attached to them.




Figure 4. The illustration above is the complete design and layout of the survey transects that will be conducted for this research. Note that the only observable portion of these surveys that will remain in the forest are the two PVC posts that mark the two ends of the 50 meters transect. These posts will have contact information and identification attached. Please refer to the key for meanings of survey types and locations.

IV. References

- 1 Sala, O.E. *et al.* (2000) Global Biodiversity Scenarios for the Year 2100. *Source Sci. New Ser.* 287, 1770–1774
- 2 Dupouey, J.L. *et al.* (2002) Irreversible Impact of past Land Use on Forest Soils and Biodiversity. *Ecology* 83, 2978–2984
- 3 Mitchell, C.E. *et al.* (2002) Effects of Historical Land Use and Forest Patch Size on Myrmecochores and Ant Communities. *Ecol. Applications* 12, 1364–1377
- 4 Bellemare, J. *et al.* (2002) Legacies of the agricultural past in the forested present: an assessment of historical land-use effects on rich mesic forests. *J. Biogeogr.* 29, 1401–1420
- 5 Ellis, E.C. and Ramankutty, N. (2008) Putting people in the map: anthropogenic biomes of the world. *Front. Ecol. Environ.* 6, 439–447
- 6 Flinn, K.M. and Vellend, M. (2005) Recovery of forest plant communities in post agricultural landscapes. *Front. Ecol. Environ.* 3, 243–250
- 7 Foster, D.R. (1992) Land-Use History (1730-1990) and Vegetation Dynamics in Central New England, USA. *J. Ecol.* 80, 753
- 8 Flinn, K.M. *et al.* (2004) Land-use history and forest herb diversity in Tomkins County, New York, USA. In *Forest Biodiversity: Lessons from History from Conservation* (Honnay O., Verheyen K., Bossuyt B., H. M., ed), pp. 81–85, CABI Pub. in association with the International Union of Forestry Research Organizations (IUFRO)
- 9 Grashof-Bokdam, C.J. and Geertsema, W. (1998) The effect of isolation and history on colonization patterns of plant species in secondary woodland. *J. Biogeogr.* 25, 837–846
- 10 Motzkin, G. *et al.* (1996) Controlling Site to Evaluate History : Vegetation Patterns of a New England Sand Plain. *Ecol. Monogr.* 66, 345–365
- 11 Griffiths, G.R. and McGee, G.G. (2018) Lack of herbaceous layer community recovery in postagricultural forests across three physiographic regions of New York. *J. Torrey Bot. Soc.* 145, 1–20
- 12 Beattie, A.J. and Culver, D.C. (1981) The Guild of Myrmecochores in the Herbaceous Flora of West Virginia Forests. *Source Ecol. Ecol.* 62, 107–115
- 13 Handel, S.N. *et al.* (1981) Ants Disperse a Majority of Herbs in a Mesic Forest Community in New York State Ants disperse a majority of herbs in a community mesic forest in New York State. *Source Bull. Torrey Bot. Club* 108, 430–437
- 14 Handel, S.N. *et al.* (1990) Seed Dispersal by Ants Thousands of plant species rely on ants to disperse their seeds. With special food lures and other adaptations, a plant can induce the insects to carry away its seeds without harming them. *Source Sci. Am.* 263, 76–83
- 15 Ness, J.H. and Morin, D.F. (2008) Forest edges and landscape history shape interactions between plants, seed-dispersing ants and seed predators. *Biol. Conserv.* 141, 838–847
- 16 Canner, J.E. *et al.* (2012) Redispersal of seeds by a keystone ant augments the spread of common wildflowers. *Acta Oecologica* 40, 31–39
- 17 Ness, J.H. *et al.* (2009) Uncommon specialization in a mutualism between a temperate herbaceous plant guild and an ant: Are Aphaenogaster ants keystone mutualists? *Oikos* 118, 1793–1804
- 18 DeMarco, B.B. and Cognato, A.I. (2016) A multiple-gene phylogeny reveals polyphyly among eastern North American Aphaenogaster species (Hymenoptera: Formicidae). *Zool. Scr.* 45, 512–520
- 19 Ellison, A.M. (2012) *A field guide to the ants of New England*, Yale University Press.